

(19) World Intellectual Property Organization
International Bureau




(43) International Publication Date
12 April 2001 (12.04.2001)

PCT


(10) International Publication Number
WO 01/24645 A1

- (51) International Patent Classification⁷: A23L 1/30, A61K 31/20 (74) Agent: BECKER, KURIG, STRAUS; Bavariastrasse 7, 80336 München (DE).
- (21) International Application Number: PCT/EP00/08995 (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (22) International Filing Date:
14 September 2000 (14.09.2000)
- (25) Filing Language: English
- (26) Publication Language: English (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- (30) Priority Data:
9923738.0 7 October 1999 (07.10.1999) GB
- (71) Applicant (*for all designated States except US*): SOCI-
ETE DES PRODUITS NESTLE S.A. [CH/CH]; P.O. Box
353, CH-1800 Vevey (CH).
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): BERGER, Alvin
[US/CH]; Nestlé Research Center, Vers-Chez-Les-Blanc,
CH-1000 Lausanne 26 (CH). CROZIER, Gayle [CA/CH];
Nestlé Research Center, Vers-Chez-Les-Blanc, CH-1000
Lausanne 26 (CH).
- Published:
— With international search report.
— Before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments.
- For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*


(54) Title: NUTRITIONAL COMPOSITION



Anandamide



2-AG



Oleamide

(57) Abstract: A nutritional or therapeutic composition for oral administration which comprises a naturally occurring precursor that is metabolised to a compound having anandamide activity for use as a medicament or nutritive product. In addition the invention includes a method of production of the composition, use of the composition in the manufacture of a nutritional composition for the treatment or prevention of a behavioural disorder; and a method of treatment or prevention of a behavioural disorder which comprises administering an effective amount of the composition. In a preferred embodiment the composition comprises a triacylglycerol having palmitate and arachidonate attached to its backbone wherein arachidonate is at the *sn*-1 and *sn*-2 positions.



WO 01/24645 A1

Nutritional Composition

5 The present invention relates to a nutritional or therapeutic composition for oral administration which comprises a naturally occurring precursor that is metabolised to a compound having anandamide activity for use as a medicament or nutritional product, a method of production of the composition, use of the composition in the manufacture of a nutritional or therapeutic composition for the treatment or prevention of a behavioural disorder; and a method of treatment or prevention of a behavioural disorder which
10 comprises administering an effective amount of the composition.

Within the context of this specification the word "comprises" is taken to mean "includes, among other things". It is not intended to be construed as "consists of only".

15 Standard nomenclature for fatty acid compounds is used. For example, the number of carbon atoms and number and position of double bonds is typified by "20:4(5,8,11,14)" for arachidonic acid: the number preceding the colon is the total number of carbon atoms, the number immediately following the colon is the number of double bonds, and the numbers in parentheses are the position of the double bonds, starting from the end of
20 the chain bearing the carboxylic acid group. In all compounds referred to in this manner, except where otherwise indicated, all double bonds are cis.

Standard nomenclature for classes of fatty acid compounds is used indicating the location of the double bond closest to the methyl end group, typified by "n-3" or "n-6":
25 the number following the dash denotes the position of the double bond closest to the methyl end of the molecule, counting from the methyl end. Thus, arachidonic acid is in the n-6 class, as is linoleic acid (18:2(9,12)), whereas eicosapentaenoic acid (20:5(5,8,11,14,17)) is in the n-3 class. This nomenclature is equivalent to "omega or ω " nomenclature in the literature, " ω " and "n" being interchangeable.

Anandamide (also referred to as *N*-arachidonylethanolamine) is an example of an *N*-acyl ethanolamine (hereinafter referred to as NAE). Both NAEs and *N*-acyl amines (hereinafter referred to as NAAs), an example of the latter of which is oleamide, are naturally occurring in the human body. They have been found in the hippocampus, striatum, cerebellum, spleen, heart, plasma and cerebral spinal fluid as well as in human milk.

The term "anandamide activity" is used within the context of this specification to mean an activity selected from the group which comprises an activity attributed to the drug 9-tetrahydrocannabinol (THC), as well as affects specific to anandamide and 1- and 2-monoarachidonylglycerol isomers (hereafter denoted AG), and unique from THC. It has been suggested that anandamide and AG activities are typically, but not necessarily, mediated by binding to the receptor class, known as CB1 and CB2 receptors. These anandamide activities include, but are not limited to: antinociception, catalepsy and inhibition of locomotor activity *in vivo* and displacement of 9-tetrahydrocannabinol (THC), inhibition of adenylate cyclase, inhibition of calcium channels, activation of phospholipase A₂, release of intracellular calcium *in vitro* and inhibition of twitch response *ex vivo*.

20

The term anandamide, as used within the context of this specification, refers to an NAE, NAA or MAG having anandamide activity (as defined above). Accepted scientific nomenclature will be used in this specification when reference is made to specific acyl moieties of an NAE, NAA or MAG.

25

It is well known that pharmaceutical compounds have wide application for their calming effects and they may be used in the treatment of patients suffering from conditions such as hypertension, glaucoma, insomnia, pain, inflammation, migraine headaches, convulsions, loss of appetite, nausea, cramps, diarrhoea, asthma, nervousness, aggressive behaviour, excessive timidity, inability to sleep, catalepsy, low mood, depression, gut upsets, or spasms, poor motor control, tics, excessive stress, spasticity or

30

multiple sclerosis. However, a number of these compounds are not naturally occurring in nature and in view of this, patients may be reluctant to be administered them. In the light of this there is a need for the provision of new products which include naturally occurring precursors of compounds that have a nutritive or therapeutic effect, when
5 metabolised endogenously to active compounds with anandamide activity.

Furthermore, a problem with most commercially available drugs is that they give rise to side affects such as nausea, bloating, cramping, etc. Clearly there is a need for a composition which does not give rise to these side effects.

10 The method of administration of a nutritive or therapeutic compound is an important consideration. Intravenous or subcutaneous administration of drugs requires expertise, and compared to oral administration it is not as safe, convenient or acceptable to the patient. In the light of these concerns it is clear that there is a need for new nutritive or
15 therapeutic products which may be administered orally.

In addition to the problems set out above, infant formulae are generally constructed so that they resemble human milk as closely as possible, however a plurality of components in human milk are bioactive and, because of synergies among the
20 components, the inclusion of only one or a few of them may not reproduce the bioactivity of human milk. In view of this, a problem which presently faces researchers lies in the formulation of infant formulae or weaning foods which have components that are present in human milk and which have an equivalent activity to human milk. The problem is compounded in view of the fact that not all of the components in human milk
25 have been identified and there are variations in the concentration of components which are present, possibly due to variations of mother's diets.

A further problem which faces nutritionists lies in the field of pet nutrition. Whereas some pets are aggressive, others are excessively timid. Muzzles have been provided
30 which fit over the heads of aggressive animals and cover their mouths. This may not be a good solution in view of the fact that a muzzle may serve to aggravate the animal. In

the light of this, there is a need for alternative solutions for calming excessively aggressive or timid pets.

US 5874459 discloses that anandamide may act as a ligand which interacts with
5 cannabinoid receptors in the central nervous system and gut (CB1 receptors) and/or
immune cells and tissues such as spleen, thymus and lymphocytes (CB2 receptors).
Furthermore, this document indicates that interactions between anandamide and these
two types of cannabinoid receptors have been shown to induce physiological effects. It
is described that non-arachidonyl NAEs and NAAs have been shown to inhibit
10 anandamide inactivating enzyme. This inhibition has the net effect of potentiating the
effect of anandamide.

It has been suggested that a family of NAEs and NAAs as well as *sn*-1 and *sn*-2
monoarachidonyl glycerides are agonists of anandamide receptors (here anandamide
15 receptor refers to a receptor that anandamide might bind to, including CB1, CB2, non-
CB receptors) and elicit responses analogous to that elicited by anandamide. The
chemical structures of NAEs and NAAs are based on fatty acids and depending on the
specific fatty acids esterified they have been shown to have different activities. For
example, whereas anandamide interacts with both the CB1 receptor of the central
20 nervous system and the CB2 receptor of the immune system, palmitolyethanolamide
may interact with the CB2 receptor but not the CB1 receptor and has an anti-
inflammatory effect but no known neural effect.

Nature, vol 396, page 636, (1998) discloses the results of an analysis wherein NAEs and
25 2 arachidonoylglycerol (2-AG) were identified from foods including human, bovine and
goat milk and cocoa at various stages of processing. The document suggests that
anandamide (300mg/kg body weight⁻¹) and 2-arachidonyl glycerol (400mg/kg body
weight⁻¹) have bioactivity when taken *orally* in mice, however the compounds were
active only at very high concentrations relative to the concentrations normally present in
30 foods and the results obtained show that the amounts of anandamide, 2-AG and
oleamide in foods, including milk and cocoa, are several orders of magnitude below

those required if administered by mouth, to reach the blood and cause observable “central effects”. However, the document indicates that pure doses of anandamide, 2-AG and oleamide have calming effects and effects on the immune system when *injected* into animals. Calming effects are characterised by lessened activity, decreased nociception and greater propensity for sleep.

US 568955 discloses that synthetically produced polyunsaturated fatty acid amides and their derivatives are able to mimic the effect of naturally occurring anandamides in the brain and bind to the cannabinoid receptor. The compounds described exhibit physiological activity and are reported as being useful active ingredients in pharmaceutical compositions for treatment of inflammation, migraines, spasticity, glaucoma and multiple sclerosis.

Remarkably it has now been found that a composition for oral administration may be provided which includes a precursor that is metabolised endogenously to form a compound having anandamide activity. It is particularly surprising that a dietary precursor is selectively taken up by the CNS and selectively incorporated into the NAE pool to serve as a CB receptor-binding ligand. In addition, it is remarkable that a dietary precursor induces only a small change in the phospholipid acyl composition but induces a large change in the NAE composition.

The invention addresses the problems set out above.

Accordingly, in a first aspect the invention provides a nutritional or therapeutic composition for oral administration which comprises a naturally occurring precursor that is metabolised to a compound having anandamide activity for use as a medicament or nutritive product.

In a second aspect the invention provides a method of production of a nutritional or therapeutic composition for oral administration which comprises the steps of identifying, purifying or synthesising a naturally occurring precursor that is metabolised

to a compound having anandamide activity.

In a third aspect the invention provides use of a precursor which is metabolised to a compound having anandamide activity in the manufacture of a nutritional or therapeutic composition for the treatment or prevention of an anandamide-mediated ailment selected from the group which comprises hypertension, glaucoma, insomnia, pain, inflammation, migraine headaches, loss of appetite, nausea, cramps, diarrhoea, gut upsets, intestinal motility disturbances, asthma, nervousness, aggressive behaviour, excessive timidity, inability to sleep, catalepsy, low mood, depression, spasms, poor motor control, tics, excessive stress, spasticity, multiple sclerosis and vocalization, poor language acquisition, skin inflammation and excess nociception.

Vocalization is taken to mean disturbances in vocalisation and vocalization related to bonding behaviour, for example between an infant and mother. Such vocalisations are important in animal husbandry and in successful nurturing of the offspring by the mother in household pets. Further, such behaviours as chronic sustained crying in human infants may be treatable by oral administration of an embodiment of a composition according to the invention.


Oral administration of an embodiment of a composition according to the invention may also be used to treat or prevent inflammation in superficial mammal tissues (e.g., skin) by modulating levels of compounds with anandamide-like activity in these tissues.

In a forth aspect the invention provides a method of treatment of an anandamide-mediated ailment selected from the group which comprises hypertension, glaucoma, insomnia, pain, inflammation, migraine headaches, loss of appetite, nausea, cramps, diarrhoea, gut upsets, intestinal motility disturbances, asthma, nervousness, aggressive behaviour, excessive timidity, inability to sleep, catalepsy, low mood, depression, spasms, poor motor control, tics, excessive stress, spasticity, multiple sclerosis and vocalization, poor language acquisition, skin inflammation and excess nociception which comprises administering an effective amount of an embodiment of the

composition according to the invention.

Preferably the precursor that is metabolised to a compound having anandamide activity comprises a long chain polyunsaturated fatty acid (LCPUFA) or derivative thereof.

5 More preferably it comprises a compound of the general formula X:



wherein R is the alkenyl moiety of a LCPUFA of total chain length 16-28 carbon atoms with 2-6 double bonds, with the first double bond at the c-1, c-3 c6, c7, c9 c12 position, counting from the non carboxyl (methyl) part of the molecule; and where R'' is selected from -H, lower alkyl, -OH, NH₃, and NHCH₂CH₂OH, or an acid addition salt or
10 complex thereof.

More preferably the precursor comprises a plurality of the formula X. Preferably 1-3 X molecules are esterified to a glycerol backbone, in the following stereochemical configurations: *sn*-1,2,3; *sn*-1,2; *sn*-1,3; *sn*-2,3; *sn*-1; *sn*-2; *sn*-3.

15

In an alternative embodiment the LCPUFA is a polyunsaturated fatty acid of 16-28 carbon atoms with 2-6 double bonds, having methylated-, branched-, cyclic-, conjugated-, non-methylene interrupted-, epoxy-, furanoid-, hydroxyl-, allylic-, trans-, and seleno- moieties.

20

More preferably the fatty acid is selected from the group which comprises arachidonate (20:4n-6 AA), linoleate (18:3n-6), gamma linolenate (18:3n-6), dihomogamma-linolenate (20:3n-6 DGLA), adrenic acid (22:4n-6), linolenate (18:3n-3), stearidonic (18:4n-3), eicosatetraenoic (20:4n-3), eicosapentaenoate (20:5n-3), docosahexaenoate (22:6n-3 DHA), docosapentaenoate (22:5n-3 or 22:5n-6), tetracosapentaenoate (24:5n-3 or 24:5n-6), tetracosahexaenoate (24:6n-3) or the Mead acid (20:3n-9).
25

Preferably, an embodiment of a composition according to the invention includes an inhibitor of anandamide inactivating enzyme (also known as amidase). Preferably the

inhibitor is selected from the group which comprises oleate and oleamide, palmitate, palmitoylethanolamide, linoleylethanolamide, 2 palmitoylglycerol, 2-linoleylglycerol.

5 Preferably an embodiment of a composition according to the invention comprises a mixture of a saturated molecule in combination with an unsaturated precursor that is metabolised to a compound having anandamide activity. Preferably, the saturated molecule is palmitate or palmitoylethanolamide. Preferably the unsaturated precursor is arachidonic acid. This provides the advantage that the anandamide activity of the metabolite formed endogenously is potentiated by both inhibiting the breakdown of a
10 metabolite having anandamide-like activity and by the saturated NAE compound binding to the CB2 receptor.

Preferably, an embodiment of a composition according to the invention comprises a mixture of a compound which reacts with a CB receptor in combination with a
15 precursor that is metabolised to a compound having anandamide activity and an inhibitor of the amidase. This provides the advantage of synergy between the active molecules and potentiation of their effect by inhibiting the breakdown of a metabolite having anandamide-like activity.

20 Preferably, the precursor that is metabolised to a compound having anandamide activity is a free fatty acid, fatty acid ester of an alcohol, or a triacylglycerol. More preferably it is a triacylglycerol having an active fatty acid at the *sn*-1 and *sn*- 2 position. This provides the advantage that it leads to particularly effective CB receptor agonism. Most preferably, the triacylglycerol comprises both the active precursor compounds (eg
25 arachidonate) and the potentiator compounds (eg palmitate). This provides the advantage of a particularly effective mixture.

Preferably, an embodiment of a composition according to the invention comprises a structured triacylglycerol prepared by the interesterification of triacylglycerols with
30 active fatty acids so that a bioactive fatty acid is found in the *sn*-2 position of the triacylglycerol. This provides the advantage of optimising delivery of the active FA to

body tissues, particularly the brain.

Preferably an embodiment of a composition according to the invention comprises a physiologically acceptable carrier diluent or adjuvant.

5

Preferably an embodiment of a composition according to the invention comprises a combination of a naturally occurring precursor that is metabolised to a compound having anandamide activity together with a typical steroidal or non-steroidal anti-inflammatory drug (NSAID). This provides the advantage that synergy occurs since the combination has the ability to diminish inflammation via different pathways.

10

Preferably, an embodiment of a composition according to the invention comprises a precursor of a CB1 receptor agonist (e.g. anandamide) in combination with a precursor of a CB2 receptor agonist (e.g. palmitoylethanolamide). This provides the advantage that the anti-pain effect of the metabolites is about 100 times stronger than the effect provided by the metabolites of either precursor individually.

15

Embodiments of the invention will now be described in further detail with reference to the accompanying drawings in which:

20

Figure 1 shows the chemical structure of *N*-arachidonyl ethanolamine (anandamide), 2-AG, and oleamide.

Figure 2 shows the effects of oral cannabimimetic lipids on ambulation, rearing, immobility and analgesia.

25

Figure 3 shows the effects of oral administration of olive oil, anandamide, 2-AG, THC and oleamide on ambulation, rearing, immobility and analgesia. Ambulation, rearing, and immobility parameters were statistically, significantly different between the treatment groups and the control group, $p < 0.01-0.05$, ANOVA, Newman-Keuls; Only THC statistically, significantly increased analgesia.

30

Figure 4 shows the effect of olive oil, anandamide, 2-AG, THC and oleamide on body temperature. All groups were statistically, significantly different from the control group, $p < 0.01-0.05$, ANOVA, Newman-Keuls.

5

Figure 5 shows the effect of olive oil, anandamide, 2-AG, THC and oleamide on faecal output. Only the THC group was statistically, significantly different from the control.

10

Figure 6 shows the changes in piglet brain *N*-acylethanolamines following dietary fatty acid modification with a scale of 0 to 250 on the axis labelled pmols/mg piglet brain lipid extract. Bars within a group of three not denoted with a letter in common are statistically significant from one another ($p < 0.01-0.05$, ANOVA, Newman-Keuls). Adeq, adequate.

15

Figure 7 shows the changes in piglet brain *N*-acylethanolamines following dietary fatty acid modification with a scale of 0 to 70 on the axis labelled pmols/mg piglet brain lipid extract. Bars within a group of three not denoted with a letter in common are statistically significant from one another ($p < 0.01-0.05$, ANOVA, Newman-Keuls). Adeq, adequate.

20

Piglets were fed using two different kinds of adapted infant formulations supplemented with low levels of arachidonate and docosahexaenoate (approximately the same levels as found in human breast milk) and obtained from different sources (see Table 1). The levels of NAE, MAG (monoacylglycerol) and primary amides were evaluated in their brains.

25

In this study piglets were fed from birth to 18 days with diets comprising embodiments of a composition according to the invention with or without 0.5% 20:4n-6 from single cell oils and 0.4% 22:6n-3 in formula, with either low (deficient) 18:2n-6(1.6%) and 18:3n-3(0.1%), or with adequate 18:2(n-6)(15.6%) and 18:3n-3(1.5%).

30

The diet compositions are shown in table 1.

Table 1 Formulas varied in n-3 and n-6 fatty acid content

Fatty Acid	18:2n-6 – 18:3n-3 deficient			18:2n-6 - 18:3n-3 adequate				
	No LCP	Egg+ fish Oil	Single cell oil	No LCP	Egg fish oil	+ single cell oil	Sow milk	
g/100g fatty acids								
8:0	8.0	7.0	7.4	17.2	15.5	14.9		
10:0	6.7	5.9	6.5	13.5	12.6	13.0		
12:0	44.2	39.7	42.9	1.0	0.3	0.3	0.1	
14:0	17.1	15.6	16.8	0.8	0.6	0.6	2.4	
16:0	9.5	10.5	9.5	11.3	12.1	10.9	28.1	
18:0	3.4	4.0	3.5	3.2	3.5	3.3	5.6	
16:1	0.1	0.34	0.1	0.1	0.3	0.2	4.7	
18:1	8.1	10.4	9.3	33.3	33.4	35.1	32.6	
18:2n-6	1.6	3.8	1.9	15.6	16.0	16.4	20.4	
18:3n-6	-	0.6	0.1	-	0.4	0.1	0.2	
20:2n-6	-	-	-	-	-	-	0.4	
20:3n-6	-	-	-	-	-	-	0.2	
20:4n-6	-	0.1	0.4	-	0.1	0.4	0.7	
22:4n-6	-	-	-	-	-	-	0.1	
18:3n-3	0.1	0.5	0.1	1.5	1.8	1.6	2.3	
20:5n-3	-	0.1	-	-	0.1	-	0.1	
22:5n-3	-	-	-	-	-	-	0.6	
22:6n-3	-	0.3	0.3	-	0.3	0.3	0.1	

5

Changes in individual brain phospholipid classes that occurred after feeding were analysed.

10 The results showed that the addition of 20:4n-6 and 22:6n-3 to diets containing adequate levels of essential fatty acids (18:2n-6 and 18:3n-3) lead to an increase in 22:6n-3 in phosphatidyl choline; a decrease in 22:5n-6 in phosphatidyl ethanolamine; and no change in arachidonate (20:4n-6) in any of the phospholipid classes.

15 Thus, the small, unsubstantial increase seen in 22:6n-3 in phosphatidyl choline is consistent with the fact that the relevant diet had added 22:6n-3; however the lack of significant increase in arachidonate in any of the phospholipid classes examined indicates that added arachidonate is not incorporated into these phospholipid classes, but

rather is metabolised or inadequately transported to the brain.

The primary amides, oleamide and arachidonamide, and 18:3 NAE were not detected and are omitted from table 2, which shows the changes in levels of MAG and NAE expressed as pmols/mg lipid that occurred following feeding of the diets.

Table 2
Monacyl glycerols (MAG)

Group	C20:4n-6	C22:4n-6	C22:6n-3
Adequate	66.0	3.53	3.87
adequate + SCO	44.4	6.23	5.93
Sow fed	44.1	6.13	6.67

Table 3

N-acyl-ethanolamines (NAE)

Group	C16:0	C18:0	C18:1n-9	C18:2n-6
Adequate	114.87	27.90	27.00	8.57
adequate + SCO	149.93	63.87	15.97	2.90
Sow fed	95.07	3.13	1.40	9.80

Table 4

N-acyl-ethanolamines (NAE)

Group	C20:4n-6	C20:5n-3	C22:4n-6	C22:5n-3	C22:6n-3
Adequate	6.10	32.87	14.80	3.63	3.80
adequate+ SCO	23.77	172.37	23.07	33.67	36.10
Sow fed	19.97	165.63	29.30	28.00	15.77

MAG levels were not statistically significant for 20:4n-6 MAG, 22:4n-6 MAG and 22:6n-3 MAG in animals fed essential fatty acid sufficient diets (*sn*-1 and 2 isomers combined). This is an important finding because specific MAGs, such as 2-AG are

known to bind to CB receptors and have bioactivity.

In animals fed the 18:2n-6/18:3n-3 sufficient diets, supplementation with AA and DHA led to increases in 20:4n-6 NAE and 22:4n-6 NAE (22:4n-6 is the 2-carbon elongation product of AA), 22:6n-3 NAE, 20:5n-3 NAE and 22:5n-3 NAE (the latter two are retroconversion products of 22:6n-3). The levels of these NAE products were similar to that found in sow milk fed piglets. Thus, it is a remarkable feature of the invention that when sufficient essential fatty acids are provided in the diet, the supplementation of AA and DHA to levels found in breast milk, has the effect of increasing corresponding NAE products to levels found in sow milk.

The results obtained indicate that supplementation with AA and DHA to formulae having sufficient essential fatty acid had minimal effects on brain phospholipid acyl moieties. However, in striking contrast, the same level of supplementation led to a 4-fold increase in the level of 20:4n-3 NAE present, a 5.2 fold increase in 20:5n-3 NAE, and a 9.5 fold increase in 22:5n-3 and 22:6n-3 NAE.

In order to show the biological activity of the composition of the present invention on animal's behaviour the effect of dietary poly-unsaturated fatty acids with and without a CB-1 receptor antagonist on anxiolytic-like responses in mice were tested. To this end, the ELEVATED PLUS MAZE TEST was applied (adapted after Handley and Mithani (Naunyn. Schmied. Arch. Pharmacol. 327: 1-5, 1984).

For the experiments male Rj:NMRI mice, obtained from Elevage Janvier, Le Genest-Saint-Isle France and weighing 10-11 g at delivery and 33-51 grams on day 42 were used. The mice were housed 10 per cage in wire cages with bedding and normal light cycle. They received *ad libitum* quantities of bottled distilled water and purified powdered diets (7.5 g/mouse) in ceramic cups (10/group) for 42 days. The Food was maintained at -80 °C in daily aliquots under nitrogen, thawed each afternoon before administration to mice. Uneaten food was discarded daily.

The principle of the test resides in that anxiolytic agents increase the number of entries into the open and often the closed arms of the elevated Consequently, mice should want to move and explore the spaces of the open and closed arms rather than staying still in the middle).

5

Mice were given the following agents intraperitoneally 60 minutes before the Plus Maze test:

Tween 80 as placebo;

the anxiolytic agent Clobazam at a non-sedative dose for test validation; or

10 validated amounts of AM251 (Tocris Cookson LTD., UK), a CB-1 receptor antagonist, to inhibit binding of endogenous NAEs to the CB-1 receptor.

All diets contained 90% fat-free AIN93G rodent diet in powder form (Lot 9350-5, Dyets, Inc., Bethlehem, PA), 0.4% milk fat, 1.2% palm olein, 1.9% Trisun sunflower oil, 1.5% soybean oil and 2.1-5.1% medium chain triacylglycerol oil. Parts of the medium chain triacylglycerol oil were replaced with 1.1% algal oil (providing 0.5 dietary wt.-% arachidonic acid) in diet D, 1.9% fish oil (providing 0.5 dietary wt.-% docosaheptaenoic acid) in diet E, and with 1.1% algal oil and 1.9% fish oil in diets F and G. Dietary groups are summarized in the table below:

20

	Diet Code	Diet Description	Agent given before Plus Maze Test
	A	Control Diet	Tween 80, 1% distilled water solution
25	B	Control Diet	Clobazam, 32 mg/kg body weight
	C	Control Diet	AM 251, 64 mg/kg body weight
	D	Diet AA	Tween 80, 1% distilled water solution
	E	Diet DHA	Tween 80, 1% distilled water solution
	F	Diet AA+DHA	Tween 80, 1% distilled water solution
30	G	Diet AA+DHA	AM 251, 64 mg/kg body weight

Abbreviations: AA, arachidonic acid; DHA, docosaheptaenoic acid

The body weight, weight gain and the food intake of the mice was monitored throughout the experiment. These parameters were not significantly affected by ingestion of the various diets using classical one way analysis of variance (ANOVA). This indicates that differences in the behavioral tests as found can only be attributed to the components in

35

the diet that were varied, namely dietary polyunsaturated fatty acids. To assess changes in the Plus Maze test, generalized Linear Models (GLM) and the *Poisson* family were used because the obtained response data are non-normally distributed counts.

5 Number of Entries in the Closed Arms

A vs. B: Average entries were 3.6 and 6.3, respectively, and the difference is at the limit of statistical significance (p -value = 0.053). This result establishes that the anxiolytic agent Clobazam, under the present conditions, can increase closed arm entries.

10 A vs. C: There was no significant difference (p -value = 0.19).

15 A, D, E and F: overall, the p -value is 0.06. The fitted average entries are respectively 3.6, 6.4, 6.0 and 6.8. In group D, there is one mouse with 16 entries, which is unusually high. Omitting this mouse, the p -value becomes significant (0.02) and the prediction for group

 D decreases to 5.3. This result establishes that the combination of dietary AA and DHA may induce anxiolytic (Clobazam-like) effects.

20 F vs. G: Average entries are respectively 6.8 and 4.6, and the difference is at the limit of statistical significance (p -value = 0.11). This result indicates that the anxiolytic effects of the combination of dietary AA and DHA may be transduced via CB-1 receptor binding, i.e., via binding of PUFA-derived NAEs.

25 C vs. G: Average entries are respectively 2.3 and 4.6, and the difference is close to statistical significance (p -value = 0.08). This result indicates that CB-1 receptors are not the only receptors that mediate responses in the PLUS MAZE TEST, i.e., non-CB-1 receptors may partially mediate the actions of dietary PUFA and (PUFA-derived) NAEs. Additionally, the drug AM251 may not fully antagonize CB-1 receptor binding.

30

In summary, the results from the number of entries into the closed arms in the

ELEVATED PLUS MAZE TEST show that dietary AA and DHA and the combination of the two, have anxiolytic-like effects that seem to be mediated via their conversion to NAEs, and these NAEs in turn bind to CB-1 receptors located in brain regions known to induce behavioral responses in the PLUS MAZE TEST, such as the hippocampus.


5

It should be understood that various changes and modifications to the presently preferred embodiments described herein will be apparent to those skilled in the art. Such changes and modifications can be made without departing from the spirit and scope of the present invention and without diminishing its attendant advantages. It is therefore intended that such changes and modifications be covered by the appended claims.

10

Claims

1. A nutritional or therapeutic composition for oral administration which comprises a naturally occurring precursor that is metabolised to a compound having anandamide activity for use as a medicament.
2. A composition according to claim 1 wherein the precursor is a long chain polyunsaturated fatty acid (LCPUFA) or derivative thereof of the general formula X:



- wherein R is the alkenyl moiety of a LCPUFA of total chain length 16-28 carbon atoms with 2-6 double bonds, with the first double bond at the c-1, c-3 c6, c7, c9 c12 position, counting from the non carboxyl (methyl) part of the molecule; and where R'' is selected from -H, lower alkyl, -OH, NH₃, and NHCH₂CH₂OH, or an acid addition salt or complex thereof.
3. A composition according to claim 2 wherein the precursor comprises a molecule having a plurality of formula X.
4. A composition according to claim 2 or 3 wherein the precursor comprises a molecule having from 1 to 3 copies of formula X esterified to a glycerol backbone, in a stereochemical configuration selected from: *sn*-1,2,3; *sn*-1,2; *sn*-1,3; *sn*-2,3; *sn*-1; *sn*-2; *sn*-3.
5. A composition according to any of the preceding claims, wherein the precursor comprises an LCPUFA which is a polyunsaturated fatty acid of 16-28 carbon atoms having from 2 to 6 double bonds, having a methylated-, branched-, cyclic-, conjugated-, non-methylene interrupted-, epoxy-, furanoid-, hydroxyl-, allylic-, trans-, or a seleno- moiety.

6. A composition according to any preceding claim wherein the precursor comprises a fatty acid selected from the group which comprises arachidonate (20:4n-6 AA), linoleate (18:3n-6), gamma linolenate (18:3n-6), dihomogamma-
linolenate (20:3n-6 DGLA), adrenic acid (22:4n-6), linolenate (18:3n-3),
stearidonic (18:4n-3), eicosatetraenoic (20:4n-3), eicosapentaenoate (20:5n-3),
docosahexaenoate (22:6n-3DHA), docosapentaenoate (22:5n-3 or 22:5n-6),
tetracosapentaenoate (24:5n-3 or 24:5n-6), tetracosahexaenoate (24:6n-3) or the
Mead acid (20:3n-9).
7. A composition according to any preceding claim wherein the precursor comprises arachidonate (20:4n-6 AA).
8. A composition according to any preceding claim which comprises an inhibitor of
an anandamide inactivating enzyme (amidase).
9. A composition according to claim 8 wherein the inhibitor is selected from the
group which comprises oleate and oleamide, palmitate, palmitoylethanolamide,
linoleylethanolamide, 2 palmitoylglycerol, 2-linoleylglycerol.
10. A composition according to claim 8 or 9 wherein the inhibitor is palmitate or
palmitoylethanolamide.
11. A composition according to any preceding claim which comprises a
triacylglycerol having palmitate and arachidonate attached to its backbone
wherein arachidonate is at the *sn*-1 and *sn*- 2 positions.
12. A composition according to any preceding claim which comprises a compound
which reacts with a CB receptor.
13. A composition according to any preceding claim which comprises a steroidal or

non-steroidal anti-inflammatory drug (NSAID).

14. A composition according to any preceding claim which comprises a physiologically acceptable carrier, diluent or adjuvant.
- 5
15. A method of production of a nutritional or therapeutic composition for oral administration which comprises the steps of identifying, purifying or synthesising a naturally occurring precursor that is metabolised to a compound having anandamide activity.
- 10
16. Use of a composition according to any one of claims 1 to 14 in the manufacture of a nutritional or therapeutic composition for the treatment or prevention of an anandamide-mediated ailment selected from the group which comprises hypertension, glaucoma, insomnia, pain, inflammation, migraine headaches, loss of appetite, nausea, cramps, diarrhoea, gut upsets, intestinal motility disturbances, asthma, nervousness, aggressive behaviour, excessive timidity, inability to sleep, catalepsy, low mood, depression, spasms, poor motor control, tics, excessive stress, spasticity, multiple sclerosis and vocalization, poor language acquisition, skin inflammation and excess nociception.
- 15
- 20
17. A method of treatment of an anandamide-mediated ailment selected from the group which comprises hypertension, glaucoma, insomnia, pain, inflammation, migraine headaches, loss of appetite, nausea, cramps, diarrhoea, gut upsets, intestinal motility disturbances, asthma, nervousness, aggressive behaviour, excessive timidity, inability to sleep, catalepsy, low mood, depression, spasms, poor motor control, tics, excessive stress, spasticity, multiple sclerosis and vocalization, poor language acquisition, skin inflammation and excess nociception which comprises administering an effective amount of a composition according to any one of claims 1 to 14.
- 25
- 30

Fig. 1






**Anandamide****2-AG****Oleamide**

Fig. 2



Ambulation, No squares
crossed

Fig. 3






Fig. 5






Fig. 6

6/7



6/7

Fig. 7



INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 00/08995

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 A23L1/30 A61K31/20

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A23L A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, BIOSIS, FSTA, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 96 40106 A (MARTEK BIOSCIENCES CORP ;KYLE DAVID J (US); LINSERT HENRY JR (US)) 19 December 1996 (1996-12-19) claims 30-78 ---	1-4,6,7, 12,14-17
A	US 5 874 459 A (HILL WILLIAM ADAM ET AL) 23 February 1999 (1999-02-23) cited in the application the whole document ---	1
A	WO 94 12466 A (YISSUM RES DEV CO ;DEVANE WILLIAM A (US); MECHOULAM RAPHAEL (IL);) 9 June 1994 (1994-06-09) claims --- -/--	1,8-10

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

G document member of the same patent family

Date of the actual completion of the international search

12 January 2001

Date of mailing of the international search report

30.01.2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Lepretre, F

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 00/08995

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 96 37200 A (SCOTIA HOLDINGS PLC ;STORDY BARBARA JACQUELINE (GB)) 28 November 1996 (1996-11-28) page 1 -page 3, paragraph 1; claims; examples ----	1-3,6,7, 12,14-17
X	EP 0 733 360 A (SCOTIA HOLDINGS PLC) 25 September 1996 (1996-09-25) claims; examples ----	1-4,6,7, 12,14-17
X	EP 0 409 559 A (EFAMOL HOLDINGS) 23 January 1991 (1991-01-23) page 5, line 5 - line 55; claims; examples ----	1-3,6, 14-17
X	WO 94 28913 A (MARTEK BIOSCIENCES CORP) 22 December 1994 (1994-12-22) page 6, line 18 - line 28; claims ----	1-4,6,7, 12,14-17
X	EP 0 490 561 A (EFAMOL HOLDINGS) 17 June 1992 (1992-06-17) claims; examples -----	1-3,6, 14-17

INTERNATIONAL SEARCH REPORT

International application No.
PCT/EP 00/08995

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claim 17 is directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 00/08995

Patent document cited in search report		Publication date	Patent family member(s)		Publication date
WO 9640106	A	19-12-1996	AU 6252196 A		30-12-1996
			EP 0831805 A		01-04-1998
US 5874459	A	23-02-1999	US 5688825 A		18-11-1997
			AU 3229097 A		05-01-1998
			EP 1021406 A		26-07-2000
			JP 2000511540 T		05-09-2000
			WO 9745407 A		04-12-1997
WO 9412466	A	09-06-1994	IL 103932 A		18-02-1997
			AU 5733494 A		22-06-1994
			CN 1097735 A		25-01-1995
			CZ 9501361 A		15-11-1995
			DE 69328492 D		31-05-2000
			DE 69328492 T		07-09-2000
			EP 0670826 A		13-09-1995
			ES 2145118 T		01-07-2000
			HU 73177 A		28-06-1996
			JP 8504195 T		07-05-1996
			PL 309051 A		18-09-1995
			SK 70895 A		13-09-1995
			US 5618955 A		08-04-1997
			ZA 9308947 A		02-08-1994
WO 9637200	A	28-11-1996	AU 722474 B		03-08-2000
			AU 5827796 A		11-12-1996
			CA 2195979 A		28-11-1996
			EP 0774962 A		28-05-1997
			FI 970298 A		24-01-1997
			JP 10503531 T		31-03-1998
			NO 970317 A		24-01-1997
			US 6150411 A		21-11-2000
			ZA 9604215 A		04-12-1996
EP 0733360	A	25-09-1996	AT 144706 T		15-11-1996
			AU 666782 B		22-02-1996
			AU 5183093 A		09-06-1994
			AU 5232996 A		18-07-1996
			CA 2109777 A		27-05-1994
			CN 1104494 A		05-07-1995
			DE 69305723 D		05-12-1996
			DE 69305723 T		03-04-1997
			DK 599576 T		25-11-1996
			EP 0599576 A		01-06-1994
			ES 2093935 T		01-01-1997
			GR 3021692 T		28-02-1997
			HK 114297 A		29-08-1997
			JP 6199663 A		19-07-1994
			NO 934266 A		27-05-1994
			NZ 250265 A		24-06-1997
			RU 2123844 C		27-12-1998
			SG 47838 A		17-04-1998
			US 5516800 A		14-05-1996
			ZA 9308835 A		02-08-1994
EP 0409559	A	23-01-1991	AT 116849 T		15-01-1995
			AU 625705 B		16-07-1992
			AU 5911590 A		24-01-1991

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 00/08995

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0409559 A		CA 2021000 A	22-01-1991
		DE 69015910 D	23-02-1995
		DE 69015910 T	08-06-1995
		DK 409559 T	27-03-1995
		ES 2066134 T	01-03-1995
		GR 3015722 T	31-07-1995
		HK 109395 A	14-07-1995
		IE 64910 B	20-09-1995
		JP 3066616 A	22-03-1991
		NZ 234528 A	24-06-1997
		ZA 9005437 A	24-04-1991
WO 9428913 A	22-12-1994	AU 693450 B	02-07-1998
		AU 6963594 A	03-01-1995
		EP 0707487 A	24-04-1996
		JP 8511533 T	03-12-1996
EP 0490561 A	17-06-1992	AT 135535 T	15-04-1996
		AU 652785 B	08-09-1994
		AU 8839791 A	11-06-1992
		CA 2056957 A	08-06-1992
		DE 69118130 D	25-04-1996
		DE 69118130 T	02-10-1996
		DK 490561 T	15-04-1996
		EP 0682878 A	22-11-1995
		ES 2084787 T	16-05-1996
		GR 3020173 T	30-09-1996
		IE 72221 B	09-04-1997
		JP 4290821 A	15-10-1992
		NZ 240841 A	27-07-1997
		US 5922345 A	13-07-1999
		ZA 9109571 A	25-11-1992